



doi • 10.5578/tt.67689
Tuberk Toraks 2018;66(4):297-303
Geliş Tarihi/Received: 12.10.2018 • Kabul Ediliş Tarihi/Accepted: 19.11.2018

Sleep disturbances in patients with lung cancer in Turkey

Yılmaz BÜLBÜL¹
Tevfik ÖZLÜ¹
Sibel ARINÇ²
Berna AKINCI ÖZYÜREK³
Hülya GÜNBATAR⁴
Ayşegül ŞENTÜRK³
Ayşe BAHADIR⁵
Melike ÖZÇELİK⁶
Ufuk YILMAZ⁷
Makbule Ö. AKBAY²
Leyla SAĞLAM⁸
Talat KILIÇ⁹
Gamze KIRKIL¹⁰
Neslihan ÖZÇELİK¹
Dursun ALİZOROĞLU⁷
Serap ARGUN BARIŞ¹¹
Durdu M. YAVŞAN¹²
Hadice S. ŞEN¹³
Serdar BERK¹⁴
Murat ACAT¹⁵
Gülfidan ÇAKMAK¹⁶
Perran F. YUMUK¹⁷
Yavuz Selim İNTEPE¹⁸
Ümran TORU¹⁹
Sibel ÖKTEM AYIK²⁰
İlknur BAŞYİĞİT¹¹
Sibel ÖZKURT²¹
Levent C. MUTLU²²

¹ Department of Chest Diseases, Farabi Hospital, Faculty of Medicine, Karadeniz Technical University, Trabzon, Turkey

¹ Karadeniz Teknik Üniversitesi Tıp Fakültesi Farabi Hastanesi, Göğüs Hastalıkları Anabilim Dalı, Trabzon, Türkiye

² Clinic of Chest Diseases, Istanbul Sureyyapasa Chest Diseases and Chest Surgery Training and Research Hospital, Istanbul, Turkey

² İstanbul Süreyyapaşa Göğüs Hastalıkları ve Göğüs Cerrahisi Eğitim ve Araştırma Hastanesi, Göğüs Hastalıkları Kliniği, İstanbul, Türkiye

³ Clinic of Chest Diseases, Ankara Atatürk Chest Diseases and Chest Surgery Training and Research Hospital, Ankara, Turkey

³ Ankara Atatürk Göğüs Hastalıkları ve Göğüs Cerrahisi Eğitim ve Araştırma Hastanesi, Göğüs Hastalıkları Kliniği, Ankara, Türkiye

⁴ Department of Chest Diseases, Faculty of Medicine, Yuzuncu Yil University, Van, Turkey

⁴ Yüzüncü Yıl Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Van, Türkiye

⁵ Clinic of Chest Diseases, Istanbul Yedikule Chest Diseases and Chest Surgery Training and Research Hospital, Istanbul, Turkey

⁵ İstanbul Yedikule Göğüs Hastalıkları ve Göğüs Cerrahisi Eğitim ve Araştırma Hastanesi, Göğüs Hastalıkları Kliniği, İstanbul, Türkiye

⁶ Division of Medical Oncology, Istanbul Lutfi Kırdar Kartal Training and Research Hospital, Istanbul, Turkey

⁶ İstanbul Dr. Lütfi Kırdar Kartal Eğitim ve Araştırma Hastanesi, Tıbbi Onkoloji Bölümü, İstanbul, Türkiye

⁷ Clinic of Chest Diseases, Izmir Dr. Suat Seren Chest Diseases and Chest Surgery Training and Research Hospital, Izmir, Turkey

⁷ İzmir Dr. Suat Seren Göğüs Hastalıkları ve Cerrahisi Eğitim ve Araştırma Hastanesi, Göğüs Hastalıkları Kliniği, İstanbul, Türkiye

⁸ Department of Chest Diseases, Faculty of Medicine, Ataturk University, Erzurum, Turkey

⁸ Atatürk Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Erzurum, Türkiye

Yazışma Adresi (Address for Correspondence)

Dr. Yılmaz BÜLBÜL

Karadeniz Teknik Üniversitesi Tıp Fakültesi Farabi Hastanesi,
Göğüs Hastalıkları Anabilim Dalı, TRABZON - TÜRKİYE

e-mail: bulbulyilmaz@yahoo.com

Zehra YAŞAR²³
Hıdır ESME²⁴
Mehmet M. EROL²⁵
Özlem ORUÇ²
Yurdanur ERDOĞAN³
Selvi ASKER⁴
Arife ULAŞ²⁶
Serhat EROL⁷
Buğra KERGET⁸
Ahmet Emin ERBAYCU⁷
Turgut TEKE¹²
Mehmet BEŞİROĞLU¹⁷
Hüseyin CAN²⁷
Ayşe DALLI²⁰
Fahrettin TALAY²³

- ⁹ Department of Chest Diseases, Turgut Ozal Medical Center, Inonu University, Malatya, Turkey
- ⁹ *İnönü Üniversitesi Turgut Özal Tıp Merkezi, Göğüs Hastalıkları Anabilim Dalı, Malatya, Türkiye*
- ¹⁰ Department of Chest Diseases, Faculty of Medicine, Fırat University, Elazığ, Turkey
- ¹⁰ *Fırat Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Elazığ, Türkiye*
- ¹¹ Department of Chest Diseases, Faculty of Medicine, Kocaeli University, Kocaeli, Turkey
- ¹¹ *Kocaeli Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Kocaeli, Türkiye*
- ¹² Department of Chest Diseases, Faculty of Medicine, Necmettin Erbakan University, Konya, Turkey
- ¹² *Necmettin Erbakan Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Konya, Türkiye*
- ¹³ Department of Chest Diseases, Faculty of Medicine, Dicle University, Diyarbakır, Turkey
- ¹³ *Dicle Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Diyarbakır, Türkiye*
- ¹⁴ Department of Chest Diseases, Faculty of Medicine, Cumhuriyet University, Sivas, Turkey
- ¹⁴ *Cumhuriyet Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Sivas, Türkiye*
- ¹⁵ Department of Chest Diseases, Faculty of Medicine, Karabük University, Karabük, Turkey
- ¹⁵ *Karabük Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Karabük, Türkiye*
- ¹⁶ Clinic of Chest Diseases, Istanbul Haseki Training and Research Hospital, Istanbul, Turkey
- ¹⁶ *İstanbul Haseki Eğitim ve Araştırma Hastanesi, Göğüs Hastalıkları Kliniği, İstanbul, Türkiye*
- ¹⁷ Division of Medical Oncology, Faculty of Medicine, Marmara University, Istanbul, Turkey
- ¹⁷ *Marmara Üniversitesi Tıp Fakültesi, Tıbbi Onkoloji Bölümü, İstanbul, Türkiye*
- ¹⁸ Department of Chest Diseases, Faculty of Medicine Bozok University, Yozgat, Turkey
- ¹⁸ *Bozok Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Yozgat, Türkiye*
- ¹⁹ Department of Chest Diseases, Faculty of Medicine, Dumlupınar University, Kutahya, Turkey
- ¹⁹ *Dumlupınar Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Kütahya, Türkiye*
- ²⁰ Department of Chest Diseases, Faculty of Medicine, Izmir Katip Celebi University, Izmir, Turkey
- ²⁰ *İzmir Katip Çelebi Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, İzmir, Türkiye*
- ²¹ Department of Chest Diseases, Faculty of Medicine, Pamukkale University, Denizli, Turkey
- ²¹ *Pamukkale Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Denizli, Türkiye*
- ²² Department of Chest Diseases, Faculty of Medicine, Namık Kemal University, Tekirdağ, Turkey
- ²² *Namık Kemal Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Tekirdağ, Türkiye*
- ²³ Department of Chest Diseases, Faculty of Medicine, Bolu Abant İzzet Baysal University, Bolu, Turkey
- ²³ *Bolu Abant İzzet Baysal Üniversitesi Tıp Fakültesi, Göğüs Hastalıkları Anabilim Dalı, Bolu, Türkiye*

- ²⁴ Clinic of Thoracic Surgery, Meram Training and Research Hospital, Konya, Turkey
- ²⁴ Meram Eğitim ve Araştırma Hastanesi, Göğüs Cerrahisi Kliniği, Konya, Türkiye
- ²⁵ Department of Thoracic Surgery, Faculty of Medicine, Uludağ University, Bursa, Turkey
- ²⁵ Uludağ Üniversitesi Tıp Fakültesi, Göğüs Cerrahisi Anabilim Dalı, Bursa, Türkiye
- ²⁶ Clinic of Medical Oncology, Atatürk Education and Research Hospital, Ankara, Turkey
- ²⁶ Atatürk Eğitim ve Araştırma Hastanesi, Medikal Onkoloji Kliniği, Ankara, Türkiye
- ²⁷ Department of Family Medicine, Faculty of Medicine, Izmir Katip Celebi University, Izmir, Turkey
- ²⁷ İzmir Katip Çelebi Üniversitesi Tıp Fakültesi, Aile Hekimliği Anabilim Dalı, İzmir, Türkiye

SUMMARY

Sleep disturbances in patients with lung cancer in Turkey

Introduction: Sleep quality is known to be associated with the distressing symptoms of cancer. The purpose of this study was to analyze the impact of cancer symptoms on insomnia and the prevalence of sleep-related problems reported by the patients with lung cancer in Turkey.

Materials and Methods: Assessment of Palliative Care in Lung Cancer in Turkey (ASPECT) study, a prospective multicenter study conducted in Turkey with the participation of 26 centers and included all patients with lung cancer, was re-evaluated in terms of sleep problems, insomnia and possible association with the cancer symptoms. Demographic characteristics of patients and information about disease were recorded for each patient by physicians via face-to-face interviews, and using hospital records. Patients who have difficulty initiating or maintaining sleep (DIMS) is associated with daytime sleepiness/fatigue were diagnosed as having insomnia. Daytime sleepiness, fatigue and lung cancer symptoms were recorded and graded using the Edmonton Symptom Assessment Scale.

Results: Among 1245 cases, 48.4% reported DIMS, 60.8% reported daytime sleepiness and 82.1% reported fatigue. The prevalence of insomnia was 44.7%. Female gender, patients with stage 3-4 disease, patients with metastases, with comorbidities, and with weight loss > 5 kg had higher rates of insomnia. Also, patients with insomnia had significantly higher rates of pain, nausea, dyspnea, and anxiety. Multivariate logistic regression analysis showed that patients with moderate to severe pain and dyspnea and severe anxiety had 2-3 times higher rates of insomnia.

Conclusion: In conclusion, our results showed a clear association between sleep disturbances and cancer symptoms. Because of that, adequate symptom control is essential to maintain sleep quality in patients with lung cancer.

Key words: Lung cancer; sleep disturbance; insomnia; symptom; pain; dyspnea

ÖZET

Türkiye’de akciğer kanseri hastalarında uyku bozuklukları

Giriş: Uyku kalitesinin, kanser semptomlarının şiddetiyle ilişkili olduğu bilinmektedir. Bu çalışmada, Türkiye’de akciğer kanseri hastalarında uyku ile ilişkili sorunların prevalansı ve kanser semptomlarının insomnia üzerine etkisinin araştırılması amaçlandı.

Materyal ve Metod: Türkiye’de 26 merkezin katılımıyla gerçekleştirilen, akciğer kanserli olguların dahil edildiği çok merkezli bir çalışma olan ASPECT çalışması verileri, bu hastalarda görülen uyku sorunları, insomni ve bunların kanser semptomları ile ilişkisi yönüyle yeniden değerlendirildi. Hastaların demografik özellikleri ve hastalıkları hakkında bilgi, hasta ile yüz yüze görüşülerek ve hastane kayıtları aracılığıyla derlendi. Uykuyu başlatma ve sürdürme zorluğu (DIMS) ile gündüz artmış uyku hali veya yorgunluk tanımlayan olgular insomni olarak değerlendirildi. Gündüz uyku hali, yorgunluk ve akciğer kanseri semptomları Edmonton Semptom Değerlendirme Skalası kullanılarak kaydedildi.

Bulgular: Katılan 1245 olgunun, %48.4’ünde DIMS, %60.8’inde gündüz uyku hali ve %82.1’inde yorgunluk mevcuttu. Insomni prevalansı %44.7 olarak bulundu. Kadın cinsiyet, evre 3-4 hastalık, metastatik hastalık, komorbid hastalıklar ve kilo kaybı > 5 kg olan olgularda insomnia oranı anlamlı olarak daha yüksekti. Diğer yandan insomnisi olan hastalarda ağrı, bulantı, dispne ve anksiyete semptomları anlamlı düzeyde yüksek bulundu. Multivariate lojistik regresyon analizi, orta ileri şiddetli ağrısı ve dispnesi olan, ayrıca ciddi anksiyetesi olan hastalarda insomni sıklığının 2-3 kat fazla olduğunu ortaya koydu.

Sonuç: Çalışmamız, kanser semptomları ile uyku kalitesi arasında yakın bir ilişki olduğunu ortaya koymuştur. Bu nedenle, akciğer kanserli hastalarda kaliteli bir uyku için, kanser semptomların yeterince kontrol edilmesi gereklidir.

Anahtar kelimeler: Akciğer kanseri; uyku bozukluğu; insomni; semptom; ağrı; dispne

INTRODUCTION

Sleep disturbances, such as difficulty falling asleep and maintaining sleep, early awakening, and excessive daytime sleepiness, are common in patients with cancer. Of the cancer patients, 30-75% report sleep problems, which is a rate about two times as high as in the general population (1). Specifically, in patients with lung cancer, 52% of newly diagnosed lung cancer patients were found to have insomnia (2). Sleep quality is known to be associated with the distressing symptoms of cancer (3). Inadequately controlled or uncontrolled cancer symptoms lead to disturbed and fragmented sleep which may contribute to poor quality of life and mood disorders, particularly depression (1).

Several studies have reported that there is a high prevalence of uncontrolled symptoms in patients with lung cancer (4,5). In a previous study, we have also shown that palliative treatment of lung cancer symptoms are mostly inadequate, and large number of patients continue to suffer from uncontrolled symptoms and unmet needs (6). Despite exiting data on association of sleep quality and symptoms in other cancer types, few studies, with limited number of patients, focused on this issue in lung cancer. We re-evaluated the results of this study to analyze the impact of cancer symptoms on insomnia and the prevalence of sleep-related problems reported by the patients with lung cancer in Turkey.

MATERIALS and METHODS

Study Design

Assessment of Palliative Care in Lung Cancer in Turkey (ASPECT) study is a prospective multicenter study conducted in Turkey (between March 2014-September 2014) with the participation of 26 centers, including all patients with lung cancer, who agreed to participate in the study (6). The study was approved by the local ethics committee and written informed consent was obtained from all patients. The results of ASPECT study were re-evaluated in terms of sleep disturbances and possible association with the cancer symptoms.

Patients and Setting

The study included all consecutive patients with lung cancer, whom disease was objectively confirmed histopathologically. Demographic characteristics of patients and information about disease were recorded for each patient by physicians via face-to-face interviews, and using hospital records.

Patients were questioned regarding whether or not they had difficulty initiating or maintaining sleep (DIMS): "Do you frequently have difficulty in falling asleep at night or waking up frequently during the night or getting back to sleep after waking during the night?" Patients with DIMS also expressed subjective reasons for DIMS as open ended questions. Daytime sleepiness, fatigue and lung cancer symptoms were recorded and graded using the Edmonton Symptom Assessment Scale (ESAS) (7). The ESAS symptom scores were categorized by severity as follows: none= 0; mild= 1-3; moderate= 4-6; and severe= 7-10. Patients who have DIMS and associated daytime sleepiness/fatigue were diagnosed as having insomnia.

Statistical Analysis

Data analysis was performed using SPSS software (Version 13.01; SPSS Inc., Chicago, IL, USA). The chi-square test was used to compare categorical variables. The parametric Student t-test was used for comparing mean or median values of normally distributed data, and the nonparametric Mann-Whitney U test was used to compare data that was not normally distributed. Demography and the symptoms that were potential predictors of insomnia were analyzed using logistic regression and multivariate logistic regression analysis was used as a stepwise descending method from predictive factors with a significance ≤ 0.05 in the univariate analysis.

RESULTS

Among 1245 cases (88.7% man) included, 48.4% (590/1218 pts) reported DIMS, 60.8% (742/1221 pts) reported daytime sleepiness and 82.1% (1002/1220) reported fatigue. The most frequent subjective reasons, reported by the patients, for DIMS were pain and dyspnea (Table 1).

The prevalence of patients with insomnia was 44.7% (541/1211 pts). Among demographic parameters,

Table 1. Subjective reasons for difficulty initiating or maintaining sleep cited by the patients with lung cancer

Reasons (590 pts)	No (%)
Pain	220 (37.3)
Dyspnea	164 (27.8)
Anxiety/stress/depression	95 (16.1)
Nocturia	70 (11.8)
Cough	33 (5.6)
Other	54 (9.1)

Table 2. Demographic parameters according to the patients with/without insomnia

Variables	Patients without insomnia No (%)	Patients with insomnia No (%)	p
Age (mean ± SD)	61.82 ± 9.45	61.76 ± 9.34	0.920
Gender			
-Man	610 (91.0)	467 (86.3)	0.009
-Woman	60 (9.0)	74 (13.7)	
BMI (mean ± SD)	24.80 ± 4.50	24.48 ± 4.63	0.119
Histopathologic type			
-Small	123 (18.4)	111 (20.5)	0.344
-NSCLC	547 (81.6)	430 (79.5)	
Time from diagnosis (mean ± SD, months)	8.99 ± 14.47	9.13 ± 13.27	0.794
Comorbidity			
-No	387 (58.2)	263 (48.7)	0.001
-Yes	278 (41.8)	277 (51.3)	
Family cancer history			
-No	468 (70.6)	347 (65.6)	0.066
-Yes	195 (29.4)	182 (34.4)	
Weight loss > 5 kg			
-No	381 (57.2)	231 (42.8)	< 0.001
-Yes	285 (42.8)	309 (57.2)	
Disease stages			
-Stage 1-2	140 (21.5)	58 (11.0)	< 0.001
-Stage 3-4	510 (78.5)	469 (89.0)	
Metastases			
-No	357 (53.7)	189 (35.1)	< 0.001
-Yes	308 (46.3)	350 (64.9)	

female patients, patients with stage 3 and 4 disease, patients with metastases, with comorbidities, and with weight loss more than 5 kg had higher rates of insomnia (Table 2). Also, lung cancer symptoms were clearly associated with insomnia. Patients with insomnia had significantly higher rates of pain, nausea, dyspnea, and anxiety (Table 3). Multivariate logistic regression analysis showed a clear association between insomnia and symptoms: pain, dyspnea and anxiety. Especially patients with moderate to severe pain and dyspnea and severe anxiety had 2 to 3 times higher rates of insomnia (Table 4).

DISCUSSION

Insomnia is a sleep disorder characterized by repeated difficulty initiating sleep, difficulty maintaining sleep, or waking up to early which results in daytime impairment despite adequate time and opportunity for sleep (8). Estimations of the insomnia prevalence depend on the criteria used to define insomnia; however a general consensus has developed from population-based studies that approximately 6-10% of patients have insomnia (9). The results of this study showed that a

Table 3. Distribution of symptoms according to the patients with/without insomnia

Symptoms	Patients without insomnia No (%)	Patients with insomnia No (%)	p
Pain			
-No	295 (44.0%)	122 (22.6%)	< 0.001
-Yes	375 (56.0%)	417 (77.4%)	
Nausea			
-No	418 (62.5%)	244 (45.3%)	< 0.001
-Yes	251 (37.5%)	295 (54.7%)	
Dyspnea			
-No	272 (40.7%)	100 (18.9%)	< 0.001
-Yes	397 (59.3%)	439 (81.4%)	
Depression			
-No	405 (60.5%)	186 (34.5%)	< 0.001
-Yes	264 (39.5%)	353 (65.5%)	
Anxiety			
-No	358 (53.4%)	154 (28.6%)	< 0.001
-Yes	312 (46.6%)	385 (71.4%)	

Table 4. Multivariate logistic regression analysis of demography and symptoms predicting insomnia

Variables*	OR	95.0% CI	p
Gender	1.273	0.829 - 1.954	0.270
Comorbidity	1.176	0.897 - 1.542	0.239
Weight loss > 5 kg	1.301	0.989 - 1.710	0.060
Stage 3-4 disease	1.311	0.852 - 2.018	0.218
Metastasis	1.319	0.966 - 1.802	0.081
Pain	REF		
Mild	0.904	0.628 - 1.303	0.589
Moderate	2.520	1.732 - 3.666	< 0.001
Severe	3.354	2.207 - 5.098	< 0.001
Nausea	REF		
Mild	0.980	0.688 - 1.397	0.913
Moderate	1.245	0.827 - 1.873	0.294
Severe	1.622	0.968 - 2.717	0.066
Dyspnea	REF		
Mild	1.354	0.929 - 1.974	0.115
Moderate	2.060	1.402 - 3.028	< 0.001
Severe	2.888	1.921 - 4.341	< 0.001
Depression	REF		
Mild	1.081	0.686 - 1.704	0.737
Moderate	1.286	0.748 - 2.213	0.363
Severe	1.597	0.838 - 3.045	0.155
Anxiety	REF		
Mild	0.979	0.624 - 1.534	0.925
Moderate	1.115	0.659 - 1.888	0.685
Severe	2.145	1.174 - 3.920	0.013

*Only variables, derived from predictive factors with a significance ≤ 0.05 in the univariate analysis, were included.

significant percentage of patients with lung cancer had poor sleep quality; insomnia (44.7%), daytime sleepiness (60.8%) and fatigue (82.1%). Ginsburg et al. reported similar results of insomnia in (52%) newly diagnosed lung cancer patients (2). Also, the prevalence of insomnia was reported to be as 31% among women patients with lung cancer (10). Chen et al. reported that about half of all lung cancer patients, receiving chemotherapy, were poor sleepers (11).

In current study, we found a significant correlation between insomnia and some demographic parameters. Female gender, patients with stage 3 and 4 disease, with metastases, with comorbidities, and patients with weight loss more than 5 kg had higher rates of insomnia. Again, we found that patients with insomnia had significantly higher rates of pain, nausea, dyspnea, and anxiety/depression. However, multivariate logistic regression analysis showed that only moderate to severe pain and dyspnea, and severe anxiety were the parameters predicting insomnia among lung cancer patients. Pain, dyspnea and anxiety/depression were also the most frequent subjective reasons of DIMS as reported by the patients. The relationship between sleep quality and the distressing symptoms of cancer is not surprising and several studies showed that some demographic parameters (including patients' gender, age, previous sleep history, cancer stage) and symptoms (due to cancer, chemotherapy and radiotherapy) were associated with the sleep problems in general cancer patients (3,12-14). The association between sleep quality and the lung cancer symptoms has also been studied in a few studies and they emphasized a relationship between sleep quality and pain, vomiting, fatigue and dyspnea and cough (11,15,16). Despite we found significantly higher rates of nausea in patients with insomnia, logistic regression analysis failed to show this association. On contrary, Akyüz et al. reported a significant correlation between sleep quality and nausea and fatigue (15). The difference between both studies might be associated with limited number of patients in the study by Akyüz et al. (15). Despite we found a relation between insomnia and distressing cancer symptoms, it is a complex process and each of these symptoms can aggravate the others, for example; unresolved symptoms, such as pain, dyspnea, fatigue etc. can lead to depression, anxiety and impaired quality of life which in turn can cause sleep disturbances and insomnia.

This study has some limitations. First of all, the results of this study primarily based on self-reported sleep disturbances with no objective measures, such as polysomnography or actigraphy. Second, we did not evaluate the effect of medications and treatment on sleep. Finally diagnosis of insomnia limited to those with DIMS and daytime sleepiness and fatigue. However in addition to daytime sleepiness and fatigue, other parameters leading daytime impairment such as attention/concentration/memory impairment, mood disturbance or irritability, motivation/energy reduction etc. were not questioned (8). Because of that, the rate of insomnia in this study might be a little bit higher than we found.

We concluded that high percentage of patients with lung cancer suffer from sleep disturbances, such as insomnia, daytime sleepiness and fatigue. Sleep disturbances in lung cancer patients were associated with cancer symptoms and especially pain, dyspnea and anxiety. Because there was a clear association between sleep disturbances and cancer symptoms, adequate symptom control is essential to maintain sleep quality in patients with lung cancer. We also conclude that, patients with lung cancer should be routinely evaluated for cancer symptoms: pain, dyspnea, and anxiety to restore sleep problems.

REFERENCES

1. Sonia Ancoli-Israel. *Sleep disturbances in cancer: a review. Sleep Med Res* 2015;6:45-9.
2. Ginsburg ML, Quirt C, Ginsburg AD, MacKillop WJ. *Psychiatric illness and psychosocial concerns of patients with newly diagnosed lung cancer. CMAJ* 1995;152:701-8.
3. McMillan SC, Tothagen C, Morgan MA. *Relationships among pain, sleep disturbances, and depressive symptoms in outpatients from a comprehensive cancer center. Oncol Nurs For* 2008;35:603-11.
4. Muers MF, Round CE. *Palliation of symptoms in non-small cell lung cancer: a study by the Yorkshire Regional Cancer Organization thoracic group. Thorax* 1993;48:339-43.
5. Di Maio M, Gridelli C, Gallo C, Manzione L, Brancaccio L, Barbera S, et al. *Prevalence and management of pain in Italian patients with advanced non-small-cell lung cancer. Br J Cancer* 2004;90:2288-96.
6. Bülbül Y, Ozlu T, Arinc S, Ozyurek BA, Gunbatar H, Senturk A, et al. *Assessment of palliative care in lung cancer in Turkey. Med Princ Pract* 2017;26:50-6.
7. Bruera E, Kuehn N, Miller MJ, Selmsler P, Macmillon K. *The Edmonton Symptom Assessment System (ESAS): a simple method for the assessment of palliative care patients. J Palliat Care* 1991;7:6-9.

8. American Academy of Sleep Medicine. *International classification of sleep disorders*. 3rd ed. IL: American Academy of Sleep Medicine, 2014.
9. Roth T. *Insomnia: definition, prevalence, etiology, and consequences*. *J Clin Sleep Med* 2007;3(Suppl 5):S7-10.
10. Sarna L. *Correlates of symptom distress in women with lung cancer*. *Cancer Pract* 1993;1:21-8.
11. Chen ML, Yu CT, Yang CH. *Sleep disturbances and quality of life in lung cancer patients undergoing chemotherapy*. *Lung Cancer* 2008;62:391-400.
12. O'Donnell JF. *Insomnia in cancer patients*. *Clin Cornerstone* 2004;6(Suppl 1):S6-14.
13. Berger AM, Parker KP, Young-McCaughan S, Mallory GA, Barsevick AM, Beck SL, et al. *Sleep wake disturbances in people with cancer and their caregivers: state of the science*. *Oncol Nurs Forum* 2005;32:E98-126.
14. Mystakidou K, Parpa E, Tsilika E, Gennatas C, Galanos A, Vlahos L. *How is sleep quality affected by the psychological and symptom distress of advanced cancer patients?* *Palliat Med* 2009;23:46-53.
15. Akyuz RC, Ugur O, Elcigil A. *Sleep quality in lung cancer patients*. *Asian Pacific J Cancer Prev* 2013;14:2909-13.
16. Gooneratne NS, Dean GE, Rogers AE, Nkwuo JE, Coyne JC, Kaiser LR. *Sleep and quality of life in long-term lung cancer survivors*. *Lung Cancer* 2007;58:403-10.